



Elements of Analysis math252-08B 2008 Ass I Solutions

1/1

① $\{x : |x^2 + x - 2| < 1\} = S \subset \mathbb{R}$. Then $|x^2 + x - 2| < 1$

$$\Leftrightarrow -1 < x^2 + x - 2 < 1$$

$$\Leftrightarrow \begin{array}{l} 0 < x^2 + x - 1 \text{ and } x^2 + x - 3 < 0 \\ 0 < (x - \alpha)(x - \beta) \text{ and } (x - \gamma)(x - \delta) < 0 \end{array}$$



Using the quadratic formula


$$\alpha = \frac{-1 - \sqrt{5}}{2}, \quad \beta = \frac{-1 + \sqrt{5}}{2}, \quad \gamma = \frac{-1 - \sqrt{12}}{2}, \quad \delta = \frac{-1 + \sqrt{12}}{2}$$

and $\gamma < \alpha < \beta < \delta$

so $S = ((-\infty, \alpha) \cup (\beta, \infty)) \cap (\gamma, \delta)$

$$= (\gamma, \alpha) \cup (\beta, \delta) = \left(\frac{-1 - \sqrt{12}}{2}, \frac{-1 - \sqrt{5}}{2} \right) \cup \left(\frac{-1 + \sqrt{5}}{2}, \frac{-1 + \sqrt{12}}{2} \right)$$

OR



$$y = |x^2 + x - 2| = |x+2||x-1|$$

so the region S corresponds

to pts x such that $y < 1$

② Let $\beta = \inf(S)$. Since b is a l.b. for S , $b \leq \beta$


If $b < \beta$ let $\epsilon = \beta - b > 0$. By (b) $\exists x \in S$ with

$$b \leq x < b + \epsilon = b + \beta - b = \beta \Rightarrow x < \beta, \text{ a contradiction (?!)}$$

since β is a l.b. Hence $b = \beta$ and $b = \inf(S)$.

$$S = \left\{ 4 - \frac{2}{n}, n = 1, 2, 3, \dots \right\}$$

$$= \left\{ 2, 3, 3\frac{1}{3}, 3\frac{1}{2}, 3\frac{2}{3}, \dots \right\}$$



Then (a) $2 = \text{glb}(S)$ since $2 \leq 4 - \frac{2}{n}$

$$\Leftrightarrow \frac{2}{n} \leq 2$$

$$\Leftrightarrow 1 \leq n \quad \text{which is true, so } 2 = \text{glb}.$$

and $2 \in S$, so it must be the $\text{glb}(S)$.

Since $4 - \frac{2}{n} < 4 \quad \forall n \in \mathbb{N}$, 4 is an u.b.

Given $\epsilon > 0$ we want n so $4 - \epsilon < 4 - \frac{2}{n} < 4$

$$\Leftrightarrow \frac{2}{n} < \epsilon \Leftrightarrow \frac{2}{\epsilon} < n$$

and we can always find such an n . Hence, by the theorem, $4 = \text{lub} S$.

Let $a_n = \frac{3n+1}{n+4}$ and let $\epsilon > 0$ be given. Working back, we

want N_ϵ so $|a_n - 3| < \epsilon$

$$\Leftrightarrow \left| \frac{3n+1}{n+4} - 3 \right| < \epsilon$$

$$\Leftrightarrow \left| \frac{3n+1 - 3n - 12}{n+4} \right| < \epsilon \Leftrightarrow \left| \frac{-11}{n+4} \right| < \epsilon$$

$$\Leftrightarrow \frac{11}{n+4} < \epsilon$$

$$\Leftrightarrow \frac{11}{\epsilon} < n+4 \Leftrightarrow n > \frac{11}{\epsilon} - 4 \quad \square$$

Let N_ϵ be any natural number with $N_\epsilon > \frac{11}{\epsilon} - 4$.

Then if $n > N_\epsilon$ we have $n > N_\epsilon > \frac{11}{\epsilon} - 4 \Rightarrow n > \frac{11}{\epsilon} - 4$

Hence \square is satisfied, hence $|a_n - 3| < \epsilon$. Thus

$\forall \epsilon > 0 \exists N_\epsilon \in \mathbb{N}$ (via $\square \rightarrow$) so $n > N_\epsilon \Rightarrow |a_n - 3| < \epsilon$.

Therefore $\lim_{n \rightarrow \infty} \frac{3n+1}{n+4} = 3$. //