

# Evaluating Jacquet's $GL(n)$ Whittaker function

Version: 4th November 2005

Kevin A. Broughan

*Department of Mathematics, University of Waikato,  
Hamilton, New Zealand  
E-mail: kab@waikato.ac.nz*

Algorithms for the explicit symbolic and numeric evaluation of Jacquet's Whittaker function for the  $GL(n, \mathbb{R})$  based generalized upper half plane for  $n \geq 2$  and an implementation for symbolic evaluation in the Mathematica package `GL(n)pack` are described. This required a careful study of the different definitions of Whittaker function which appear in the literature.

*Key Words:* K-Bessel function, Whittaker function, Jacquet Whittaker function, symbolic evaluation, quadrature, unbounded domain

MSC2000: 33C15, 22E30, 11E57, 11E76.

## 1. INTRODUCTION

## 2. DEFINITIONS

Use [4] as the standard for definitions and point out where we have found ones that differ in the literature.

$GL(n)$  means  $GL(n, \mathbb{R})$ . Always, unless otherwise noted,  $n \geq 2$ .

From [4, Definition 2.4.1], let  $b_{i,j} := ij$  when  $i + j \leq n$  and  $(n - i)(n - j)$  otherwise, and  $\nu = (\nu_1, \dots, \nu_{n-1}) \in \mathbb{C}^{n-1}$ , then the so-called **power function**  $I_\nu : \mathfrak{h}^n \rightarrow \mathbb{C}$  is defined by:

$$(1) \quad I_\nu(z) := \prod_{i=1}^{n-1} \prod_{j=1}^{n-1} y_i^{b_{i,j} \nu_j},$$

where  $z \equiv x.y$  is the **Iwasawa form**, i.e.  $z = x.y.o.d$ , where  $o$  is orthogonal,  $d$  is non-zero and in the center and

$$x.y = \begin{pmatrix} 1 & x_{1,2} & x_{1,3} & & x_{1,n} \\ & 1 & x_{2,3} & & x_{2,n} \\ & & \ddots & & \vdots \\ & & & 1 & x_{n-1,n} \\ & & & & 1 \end{pmatrix} \cdot \begin{pmatrix} y_1 y_2 \cdots y_{n-1} & & & & \\ & y_1 y_2 \cdots y_{n-2} & & & \\ & & \ddots & & \\ & & & y_1 & \\ & & & & 1 \end{pmatrix}.$$

1

and both  $x$  and  $y$  are unique. Examples of the power function in dimensions 2,3,4:

$$\begin{aligned} I_\nu(y_1) &= y_1^{\nu_1}, \\ I_\nu(y_1, y_2) &= y_1^{\nu_1+2\nu_2} y_2^{2\nu_1+\nu_2}, \\ I_\nu(y_1, y_2, y_3) &= y_1^{\nu_1+2\nu_2+3\nu_3} y_2^{2\nu_1+4\nu_2+2\nu_3} y_3^{3\nu_1+2\nu_2+\nu_3}. \end{aligned}$$

Note that Friedberg [3] and Stade [11] reverse the order of the  $y_i$ 's in the Iwasawa form.

**DEFINITION 2.1.** Let  $S = U_n(\mathbb{R})$  be the subgroup of upper triangular unipotent matrices. A function  $\psi : S \rightarrow \mathbb{C}$  which can be written in the form

$$\psi(u) = \prod_{i=1}^{n-1} e^{2\pi i m_{n-i+1} u_{i,i+1}},$$

for some  $n-1$  tuple of integers  $m = (m_1, \dots, m_{n-1})$ , is called a **character** or character of  $U_n(\mathbb{R})$ . We write  $\psi_m$  for  $\psi$ , and in case each  $m_i = 1$  write  $\psi_1$ . Note that  $\psi(a.b) = \psi(a)\psi(b)$  for  $a, b \in U_n(\mathbb{R})$  and that all characters of  $U_n(\mathbb{R})$  have this form. Note also that [4] begins with a direct order for the  $m_i$  and then reverses the order for the definition of the Jacquet Whittaker function as given here.

From [4, Definition 5.9.2], if  $n \geq 2$  and  $\nu = (\nu_1, \dots, \nu_{n-1})$ :

$$\begin{aligned} \Gamma_\nu &:= \prod_{j=1}^{n-1} \prod_{j \leq k \leq n-1} \pi^{-\frac{1}{2} - v_{j,k}} \Gamma\left(\frac{1}{2} + v_{j,k}\right) \text{ where} \\ v_{j,k} &:= \sum_{i=0}^{j-1} \frac{n\nu_{n-k+i} - 1}{2}. \end{aligned}$$

**DEFINITION 2.2.** [4, Proposition 2.3.1] The associative algebra  $D^n$  is the algebra of operators generated by real linear combinations of the operators  $D_{\alpha_1} \circ \dots \circ D_{\alpha_k}$  where each  $\alpha_i$  is an  $n \times n$  real matrix,  $D_\alpha$  is defined for smooth functions  $F$  acting on elements  $g \in GL(n, \mathbb{R})$  by

$$D_\alpha F(g) := \frac{\partial}{\partial t} F(g + tg.\alpha)|_{t=0},$$

and  $D_\alpha \circ D_\beta$  is the composition of operators. The center of this algebra is denoted  $\mathfrak{D}^n$ .

**DEFINITION 2.3.** [4, Definition 1.3.1] Let  $a, b \geq 0$ . The **Siegel set**  $\Sigma_{a,b} \subset \mathfrak{h}^n$  is the set of all  $z = x.y \in \mathfrak{h}^n$  with  $|x_{i,j}| \leq b$  for  $1 \leq i < j \leq n$  and  $y_i > a$  for  $1 \leq i \leq n-1$ .

From [4, Definition 5.4.1], for  $n \geq 2$  and  $\nu = (\nu_1, \dots, \nu_{n-1})$  and  $\psi$  a character of  $U_n(\mathbb{R})$ , a smooth function  $W : \mathfrak{h}^n \rightarrow \mathbb{C}$  is called an  $SL(n, \mathbb{Z})$ -Whittaker function of type  $\nu$  (or for short a **Whittaker function**) if it satisfies the following conditions:

- (1)  $W(uz) = \psi(u)W(z)$  for all  $u \in U_n(\mathbb{R}), z \in \mathfrak{h}^n$ ,
- (2)  $DW(z) = \lambda_D W(z)$  for all  $D \in \mathfrak{D}^n, z \in \mathfrak{h}^n$ ,
- (3)  $\int_\Omega |W(z)|^2 d^*z < \infty$  where  $\Omega$  is  $\Sigma_{\frac{\sqrt{3}}{2}, \frac{1}{2}}$  and  $d^*z$  is the left invariant quotient measure.



### 3. RELATION TO STADE'S WHITTAKER FUNCTION

The  $GL(n)$ pack function Whittaker computes a symbolic iterated integral representation of the generalized Jacquet Whittaker function  $W_{Jacquet}$  (also written  $W_J$ ) of order  $n$ , for  $n \geq 2$ , as defined above. The algorithm uses the recursive representation of the Whittaker function derived by Stade [Stade, 1990, Theorem 2.1], but his Whittaker functions are not the same as those of [4]. Let  $W_S$  and  $W_S^*$  be Stade's Whittaker and Whittaker starred functions respectively and let  $\Gamma_\nu$  represent the gamma factors for either form as given in Definition 2.1 above. Make the following additional definitions:

DEFINITION 3.1.

$$\begin{aligned} H_\nu(y) &:= I_\nu(y_{n-1}, \dots, y_1), \\ Q = Q_\nu(y) &:= H_\nu(y) \prod_{j=1}^{n-1} y_j^{-\mu_j} \text{ where,} \\ \mu_j &:= \sum_{k=1}^{n-j} r_{j,k} \text{ for } 1 \leq j \leq n-1 \text{ and where,} \\ r_{j,k} &:= \left( \sum_{i=k}^{k+j-1} \frac{n\nu_i}{2} \right) - \frac{j}{2} \text{ for } 1 \leq j \leq n-1, 1 \leq k \leq n-j. \end{aligned}$$

DEFINITION 3.2.

$$\begin{aligned} W_S(y; \nu, \psi_1) = W_{n,\nu}(y) &:= \Gamma_\nu \int_{U_n(\mathbb{R})} H_\nu(w \cdot u \cdot y) \overline{\psi_1(u)} d^*u, \\ W_S^*(y; \nu, \psi_1) = W_{n,\nu}^*(y) &:= W_S(y; \nu, \psi_1) / Q. \end{aligned}$$

The notations  $W_{n,\nu}, W_{n,\nu}^*$  are from [11]. Note that the first differs from his  $W_{n,a}$  [12].

THEOREM 3.1. *Let  $n \geq 2$  and for  $y = (y_1, \dots, y_{n-1})$  let  $y_r = (y_{n-1}, \dots, y_1)$  be the vector with coordinates reversed. Then the relationship between the two definitions of Whittaker function for  $\mathfrak{h}^n$  may be expressed by the equalities:*

$$Q_\nu(y_r) * W_S^*(y_r; \nu, \psi_1) = \Gamma_\nu * W_J(y; \nu, \psi_1) = W_J^*(y; \nu, \psi_1) = W_S(y_r; \nu, \psi_1).$$

Stade's recursive formula for the Whittaker function:

THEOREM 3.2. [11, Theorem 2.1] as amended in [12] *If  $n \geq 3$  and  $\nu \in \mathbb{C}^{n-1}$ , for  $2 \leq j \leq n-2$  let  $\lambda = (\lambda_1, \dots, \lambda_{n-3})$  where  $\lambda_{j-1} := n\nu_j / (n-2)$ , set  $u_0 = 0, 1/u_{n-1} = 0$  and  $u_{n-1}^0 = 1$ .*

$$\begin{aligned} W_S^*(y; \nu, \psi_1) &= 1 \text{ if } n = 0 \text{ or } 1, \\ W_S^*(y; \nu, \psi_1) &= 2K_{\nu - \frac{1}{2}}(2\pi y_1) \text{ if } n = 2, \\ W_S^*(y; \nu, \psi_1) &= 8 \int_0^\infty u^{\frac{3\nu_1 - 3\nu_2 - 2}{2}} K_{\frac{3\nu_1 + 3\nu_2 - 2}{2}}(2\pi \sqrt{1 + \frac{1}{u_1^2}} y_1) K_{\frac{3\nu_1 + 3\nu_2 - 2}{2}}(2\pi \sqrt{1 + u_1^2} y_2) du_1 \text{ for } n = 3, \\ W_S^*(y; \nu, \psi_1) &= 2^{2n-3} \int_{(\mathbb{R}^+)^{n-2}} \left\{ \prod_{i=1}^{n-1} u_i^{r_{i,1} - r_{i,n-i}} K_{\mu_1}(2\pi y_i \sqrt{(1 + u_{i-1}^2)(1 + 1/u_i^2)}) \right\} \\ &\quad \times W_S^*\left(\left(\frac{y_2 u_1}{u_2}, \dots, \frac{y_{n-2} u_{n-3}}{u_{n-2}}\right), (\lambda_1, \dots, \lambda_{n-3})\right) \prod_{i=1}^{n-2} \frac{du_i}{u_i} \end{aligned}$$

for  $n \geq 4$ , where the quantities  $r_{i,j}$  are defined in terms of the  $\nu_i$  in Definition 3.1 above.

#### 4. CLASSICAL WHITTAKER FUNCTIONS

Properties of classical Whittaker functions are well known. However we record them here to show the relationship between the classical and  $GL(n)$ pack functions.

Whittaker's equation [16, 8, 9] for  $W_{k,\mu}(z)$  c/- [4, p57] is given by:

$$w'' + \left(-\frac{1}{4} + \frac{k}{z} + \frac{\frac{1}{4} - \mu^2}{z^2}\right)w = 0$$

where  $\mu \in \mathbb{C}$ ,  $k \in \mathbb{R}$  and  $z \in \mathbb{C}$ .

Solutions for this equation have the integral representation [16, 8, 9]:

$$W_{k,\mu}(z) = \int_0^\infty e^{-t} t^{\mu-k-\frac{1}{2}} \left(1 + \frac{t}{z}\right)^{\mu-k-\frac{1}{2}} dt$$

for  $\Re\mu - k - \frac{1}{2} > 0$ .

Solutions also have a series representation [16, 8, 9]: Let  $\Psi(\alpha, \gamma; z)$  be the so-called **confluent hypergeometric function of the second kind** satisfying

$$\Psi(\alpha, \gamma; z) = z^{-\alpha} \left( \sum_{k=0}^n \frac{(-1)^k (\alpha)_k (1 + \alpha - \gamma)_k}{k!} z^{-k} + O\left(\frac{1}{|z|^{n+1}}\right) \right),$$

for  $|\arg z| < \pi - \delta$  for all fixed  $\delta > 0$ . Then we can write

$$W_{k,\mu}(z) = z^{\mu+\frac{1}{2}} e^{-\frac{z}{2}} \Psi\left(\frac{1}{2} - k + \mu, 2\mu + 1; z\right).$$

#### 5. EVALUATION OF THE COMPLEX ORDER FUNCTION $K_\nu(X)$ .

The expressions for Whittaker functions are given in terms of the modified Bessel function of the second order  $K_\nu(z)$ , rather than  $W_{k,\nu}(z)$ . This is traditional rather than essential, but should be useful, given the way K-Bessel functions appear in many other number-theory oriented computations. Define for  $\Re z > 0$ , and  $\nu \in \mathbb{C}$  [4, p57] or [15]:

$$K_\nu(z) = \frac{1}{2} \int_0^\infty e^{-\frac{z}{2}(t+\frac{1}{t})} t^{\nu-1} dt$$

Or for  $|\arg z| < \pi$ , [8, p204]:

$$K_\nu(z) = \sqrt{\pi} (2z)^\nu e^{-z} \Psi\left(\nu + \frac{1}{2}, 2\nu + 1; 2z\right).$$

Asymptotic values for  $\Re\nu > 0$  [9, p454] as  $z \rightarrow \infty$  with  $|\arg z| \leq 3\pi/2 - \delta$ :

$$K_\nu(z) \sim \sqrt{\frac{\pi}{2z}} e^{-z}$$

Asymptotic values for  $\Re\nu > 0$  [9, p454] as  $z \rightarrow 0$  with  $|\arg z| \leq B$ :

$$K_\nu(z) \sim \frac{\Gamma(\nu)}{2} \left(\frac{z}{2}\right)^{-\nu}$$

Integral relations [9, p254]. For  $\Re\mu > |\Re\nu|$ :

$$\int_0^\infty t^{\mu-1} K_\nu(t) dt = 2^{\mu-2} \Gamma\left(\frac{\mu+\nu}{2}\right) \Gamma\left(\frac{\mu-\nu}{2}\right),$$

and for  $\Re\nu > -\frac{1}{2}$ ,  $|\arg z| < \frac{\pi}{2}$ :

$$K_\nu(z) = \frac{\sqrt{\pi}(\frac{1}{2}z)^\nu}{\Gamma(\nu + \frac{1}{2})} \int_1^\infty e^{-zt} (t^2 - 1)^{\nu - \frac{1}{2}} dt.$$

By [5, §3.384, 9.6], If  $\Re\nu > \frac{1}{2}$  and  $p \neq 0$ :

$$\int_{-\infty}^\infty \frac{e^{-ipx}}{(x^2 + 1)^\nu} dx = \frac{2\pi}{2^\nu} \frac{|p|^\nu}{\Gamma(\nu)} W_{0, \frac{1}{2} - \nu}(2|p|).$$

The system Mathematica<sup>TM</sup> gives for  $\Re\nu > \frac{1}{2}$  and  $y \neq 0$ :

$$\int_{-\infty}^\infty \frac{e^{-2\pi i y x}}{(x^2 + 1)^\nu} dx = \frac{2\pi^\nu |y|^{\nu - \frac{1}{2}}}{\Gamma(\nu)} K_{\nu - \frac{1}{2}}(2\pi|y|),$$

where the property  $K_s(z) = K_{-s}(z)$  has been used.

It follows from the series representations for the K-Bessel and classical Whittaker functions that for all  $\nu$  and  $|\arg z| < \pi - \delta$ ,

$$\sqrt{\frac{2z}{\pi}} K_\nu(z) = W_{0, \nu}(2z).$$

Comparing this with Stade's  $W_S$  in dimension  $n=2$  the reason for the term "Whittaker" finally comes clear.

## 6. UNIFORMIZATION OF THE DEFINITIONS OF WHITTAKER FUNCTION IN DIMENSIONS 2 AND 3

**Dimension 2:** We have [4, Eqn. 5.5.4]:

$$W_J(z; \nu, \psi_m) = \frac{2|m|^{\nu - \frac{1}{2}} \pi^\nu}{\Gamma(\nu)} \sqrt{y} K_{\nu - \frac{1}{2}}(2\pi|m|y) \cdot e^{2\pi i m x}$$

$$\Gamma_\nu = |m\pi|^{-\nu} \Gamma(\nu).$$

Stade's form for dimension 2 [11] is:

$$W_S(y; \nu, \psi_1) = 2\sqrt{y} K_{\nu - \frac{1}{2}}(2\pi y).$$

**THEOREM 6.1.** *Let  $n = 2$  so  $z = x + iy$ . Assume for  $y > 0$  and  $\Re\nu > \frac{1}{2}$  that*

$$W_S(y, \nu, \psi_1) = 2\sqrt{y} K_{\nu - \frac{1}{2}}(2\pi y) \text{ and } W_J(z, \nu, \psi_m) = y^{1-\nu} \int_{-\infty}^\infty \frac{e^{-2\pi i m u y}}{(1 + u^2)^\nu} du \cdot e^{2\pi i m x}.$$

*Then*

$$W_J(z, \nu, \psi_m) = |m|^{\nu-1} \cdot \left(\frac{\pi^\nu}{\Gamma(\nu)}\right) \cdot 2\sqrt{y} K_{\nu - \frac{1}{2}}(2\pi|m|y) \cdot e^{2\pi i m x}.$$



2. It follows that

$$\begin{aligned}
W_J(Mz; \nu, \psi_{\epsilon_1, \dots, \epsilon_{n-1}}) &= \int_{U_n(\mathbb{R})} I_\nu(w.u.Mz) e^{-2\pi i(\epsilon_1 u_1 + \dots + \epsilon_{n-1} u_{n-1})} d^* u \\
&= J\left(\frac{u}{\hat{u}}\right) \int_{U_n(\mathbb{R})} I_\nu(w.M\hat{u}.z) e^{-2\pi i(|m_1|\epsilon_1 \hat{u}_1 + \dots + |m_{n-1}|\epsilon_{n-1} \hat{u}_{n-1})} d^* \hat{u} \\
&= J\left(\frac{u}{\hat{u}}\right) \int_{U_n(\mathbb{R})} I_\nu(w.M.w.w.\hat{u}.z) e^{-2\pi i(m_1 \hat{u}_1 + \dots + m_{n-1} \hat{u}_{n-1})} d^* \hat{u} \\
&= \prod_{i=1}^{n-1} \prod_{j=1}^{n-1} |m_i|^{-b_{i,j} \nu_j} J\left(\frac{u}{\hat{u}}\right) \int_{U_n(\mathbb{R})} I_\nu(\omega_n.\hat{u}.z) e^{-2\pi i(m_1 \hat{u}_1 + \dots + m_{n-1} \hat{u}_{n-1})} d^* \hat{u} \\
&= \gamma_{m,\nu} \cdot W_J(z; \nu, \psi_m) \quad (1)
\end{aligned}$$

where we have used  $w.M.w$  is a diagonal matrix with elements in the reverse order from those of  $M$  and where

$$\gamma_{\nu,m} = \frac{\prod_{i=1}^{n-1} |m_i|^{i(n-i)}}{\prod_{i=1}^{n-1} \prod_{j=1}^{n-1} |m_i|^{b_{i,j} \nu_j}}.$$

3. Next, by taking the Iwasawa form for  $z = x.y$ , commuting  $M$  and  $x$  ( $M.x = \hat{x}.M$ ), and then making the transformation  $\hat{u} = u.\hat{x}$  (which has Jacobian 1), we obtain the form

$$W_J(Mz; \nu, \psi_{\epsilon_1, \dots, \epsilon_{n-1}}) = \psi_m(x) \cdot W_J(My; \nu, \psi_{\epsilon_1, \dots, \epsilon_{n-1}}) \quad (2)$$

4. Now consider the  $n - j^{\text{th}}$  row of the matrix  $u$  with  $1 \leq j \leq n - 1$ :

$$(0, \dots, 0, 1, u_j, u_{n-j, n-j+2}, \dots, u_{n-j, n}).$$

If  $\delta_j$  is the diagonal matrix with 1's in every position except the  $n - j^{\text{th}}$  which is  $\epsilon_j$ , and we make the transformation  $\hat{u} = \delta_j.u$  with Jacobian determinant  $\epsilon_j^j$ , then, since  $w.\delta_j = \delta_{n-j+1}.w$  and  $I_\nu(\delta_{n-j+1}.z) = I_\nu(z)$ , (because  $\delta_{n-j+1}$  is orthogonal and diagonal matrices commute), we can write:

$$\begin{aligned}
W_J(My; \nu, \psi_{\epsilon_1, \dots, \epsilon_{n-1}}) &= \int_{U_n(\mathbb{R})} I_\nu(w.\delta_j.\delta_j.u.My) e^{-2\pi i(\epsilon_1 u_1 + \dots + \epsilon_{n-1} u_{n-1})} d^* u \\
&= |\epsilon_j^j| \int_{U_n(\mathbb{R})} I_\nu(\delta_{n-j+1}.w.\hat{u}.My) e^{-2\pi i(\epsilon_1 \hat{u}_1 + \dots + \hat{u}_j + \dots + \epsilon_{n-1} \hat{u}_{n-1})} d^* \hat{u} \\
&= W_J(My; \nu, \psi_{\epsilon_1, \dots, 1, \dots, \epsilon_{n-1}})
\end{aligned}$$

where the subscript 1 is in the  $j^{\text{th}}$  position.

One may do the above procedure for each  $j$  with  $1 \leq j \leq n - 1$  to obtain:

$$W_J(My; \nu, \psi_{\epsilon_1, \dots, \epsilon_{n-1}}) = W_J(My; \nu, \psi_{1, \dots, 1}) \quad (3).$$

Finally, combining the expressions (1), (2) and (3) we derive the equation

$$W_J(z; \nu, \psi_m) = c_{\nu,m} \cdot \psi_m(x) \cdot W_J(My; \nu, \psi_{1, \dots, 1})$$

where

$$\begin{aligned}
c_{\nu,m} &= \gamma_{\nu,m}^{-1} \\
&= \prod_{i=1}^{n-1} |m_i|^{\sum_{j=1}^{n-1} b_{i,j} \nu_j - i(n-i)}.
\end{aligned}$$

Here is a listing of the first  $n = 2$  thru  $n = 5$   $c_{m,\nu}$  values:

$$\begin{aligned}
c_{2,\nu} &= |m_1|^{\nu_1-1}, \\
c_{3,\nu} &= |m_1|^{\nu_1+2\nu_2-2} |m_2|^{2\nu_1+\nu_2-2}, \\
c_{4,\nu} &= |m_1|^{\nu_1+2\nu_2+3\nu_3-3} |m_2|^{2\nu_1+4\nu_2+2\nu_3-4} |m_3|^{3\nu_1+2\nu_2+\nu_3-3}, \\
c_{5,\nu} &= |m_1|^{\nu_1+2\nu_2+3\nu_3+4\nu_4-4} |m_2|^{2\nu_1+4\nu_2+6\nu_3+3\nu_4-6} \\
&\quad \times |m_3|^{3\nu_1+6\nu_2+4\nu_3+2\nu_4-6} |m_4|^{4\nu_1+3\nu_2+2\nu_3+\nu_4-4}.
\end{aligned}$$

## 8. COMPUTATION OF $W_J(Z, \nu, \psi)$ AND VALIDATION

Computation of the Whittaker function was divided into symbolic evaluation and numeric evaluation. The former is more straight forward than the latter and was able to be included in `GL(n)pack`. The numerical code uses the symbolic form as an initial step. Stade's form  $W_S^*$  was computed using his recursive reformulation Theorem 3.1, which was converted first to a single multiple integral and then, by a change of variables using the inverse hyperbolic tangent in each variable, to an integral over a cube of appropriate dimension. This has a number of decided advantages over any direct use of Jacquet's integral for numerical computation: firstly the oscillation implied by the character  $\psi_1$  is removed, and secondly the exponential decay of the K-Bessel functions at infinity assists the speed and accuracy of any quadrature application.

Stade's form was then converted into the function  $W_J$  using Theorem 3.1.

Examples of the `GL(n)pack` output are given below, in Figure 1 dimensions 2, 3, in Figure 2 dimension 4 and in Figure 3 dimension 5 [2]:

`In[114]:=`

`Whittaker[{{y1, 0}, {0, 1}}, {v1}, {1}, u][[4]]`

`Out[114]=`

$$\frac{2 \pi^{v_1} \sqrt{y_1} K[-\frac{1}{2} + v_1, 2 \pi y_1]}{\Gamma[v_1]}$$

`In[115]:=`

`Whittaker[{{y1 y2, 0, 0}, {0, y1, 0}, {0, 0, 1}}, {v1, v2}, {1, 1}, u][[4]]`

`Out[115]=`

$$\begin{aligned}
&\left( 8 \pi^{-\frac{1}{2}+3v_1+3v_2} y_1^{1+\frac{v_1}{2}-\frac{v_2}{2}} y_2^{1-\frac{v_1}{2}+\frac{v_2}{2}} \int_0^\infty u^{-1+\frac{3(v_1-v_2)}{2}} K\left[\frac{1}{2}(-2+3v_1+3v_2), 2\pi\sqrt{1+u^2} y_1\right] \right. \\
&\quad \left. K\left[\frac{1}{2}(-2+3v_1+3v_2), 2\pi\sqrt{1+\frac{1}{u^2} y_2}\right] du \right) / \\
&\left( \Gamma\left[\frac{3v_1}{2}\right] \Gamma\left[\frac{3v_2}{2}\right] \Gamma\left[\frac{1}{2}(-1+3v_1+3v_2)\right] \right)
\end{aligned}$$

FIG. 1. The `GL(n)pack` Whittaker functions in dimensions 2 and 3.

To validate the numerical computations (and thus the symbolic forms computed by `GL(n)pack`), and give some idea of their accuracy, we used a result highlighted in [12, p126], namely that if the power function is defined using some especially chosen new parameters, then the Whittaker functions are invariant under all permutations of those parameters. These permutations give rise to functional equations, which on the face of it differ from those set out in [4, Theorem 5.9.8]. These permutations were used here in a simpler manner: a permutation of an explicit set of values for the new parameters

In[116]:=

```
Whittaker[{{y1 y2 y3, 0, 0, 0}, {0, y1 y2, 0, 0}, {0, 0, y1, 0}, {0, 0, 0, 1}},
{v1, v2, v3}, {1, 1, 1}, u][[4]]
```

Out[116]=

$$\left( 64 \pi^{-2+6v_1+8v_2+6v_3} y_1^{\frac{3}{2}+v_1-v_3} y_2^2 y_3^{\frac{3}{2}-v_1+v_3} \int_0^\infty \int_0^\infty K\left[-\frac{1}{2} + 2v_2, \frac{2\pi y_2 u[1]}{u[2]}\right] K\left[-\frac{3}{2} + 2v_1 + 2v_2 + 2v_3, 2\pi y_3 \sqrt{1 + \frac{1}{u[1]^2}}\right] K\left[-\frac{3}{2} + 2v_1 + 2v_2 + 2v_3, 2\pi y_2 \sqrt{(1+u[1]^2) \left(1 + \frac{1}{u[2]^2}\right)}\right] K\left[-\frac{3}{2} + 2v_1 + 2v_2 + 2v_3, 2\pi y_1 \sqrt{1+u[2]^2}\right] (u[1] u[2])^{-1+2v_1-2v_3} du[2] du[1] \right) /$$

$$\left( \Gamma[2v_1] \Gamma[2v_2] \Gamma\left[-\frac{1}{2} + 2v_1 + 2v_2\right] \Gamma[2v_3] \Gamma\left[-\frac{1}{2} + 2v_2 + 2v_3\right] \Gamma[-1 + 2v_1 + 2v_2 + 2v_3] \right)$$

FIG. 2. The GL(n)pack Whittaker functions in dimension 4.

$a_i$  give rise to two corresponding sets of values in the original parameters  $\nu_i$ . These corresponding sets should be in or close to the domain of absolute convergence of the Whittaker function  $\Re\nu_i > 1/n$  to give convergence of the integral forms.

In more detail, set

$$H_{n,a}(y) := \prod_{j=1}^{n-1} y_j \prod_{j=1}^{n-1} y_j^{a_j}$$

where the  $(a_j)$  are  $n - 1$  complex numbers. Then set  $a_n = -a_1 - \dots - a_{n-1}$ . When defined using this power function the Whittaker function is invariant under all permutations of the  $(a_i)$ . Then define  $\nu$  in terms of  $a$  by setting  $I_\nu(y) = H_{n,a}(y_r)$  and note that the first product term in the definition of  $H$  is invariant under reversal of the order of the  $y_i$ . These relations in dimensions 3 through 5 are as follows:

$$\begin{aligned} \text{Dimension 3 : } \nu_1 &= (1 + a_1 + 2a_2)/3 \\ \nu_2 &= (1 + a_1 - a_2)/3 \\ \text{Dimension 4 : } \nu_1 &= (1 + a_1 + a_2 + 2a_3)/4 \\ \nu_2 &= (1 + a_2 - a_3)/4 \\ \nu_3 &= (1 + a_1 - a_2)/4 \\ \text{Dimension 5 : } \nu_1 &= (1 + a_1 + a_2 + a_3 + 2a_4)/5 \\ \nu_2 &= (1 + a_3 - a_4)/5 \\ \nu_3 &= (1 + a_2 - a_3)/5 \\ \nu_4 &= (1 + a_1 - a_2)/5 \end{aligned}$$

In this study, the Mathematica general adaptive quadrature routine NIntegrate was used, with the option Method set of MultiDimensional and the precision set to MachinePrecision. The processor was an Intel Pentium 4. No improvement was found using the function Compile. This is no doubt because most of the work is done by NIntegrate, which is already compiled. The values given are for the

`In[1]:= Whittaker[{{y1 y2 y3 y4, 0, 0, 0, 0}, {0, y1 y2 y3, 0, 0, 0}, {0, 0, y1 y2, 0, 0},  
{0, 0, 0, y1, 0}, {0, 0, 0, 0, 1}}, {v1, v2, v3, v4}, {1, 1, 1, 1}, u][[4]]`

$$\begin{aligned}
\text{Out[1]} = & \left( 1024 \pi^5 {}^{(-1+2v_1+3v_2+3v_3+2v_4)} \right. \\
& y_1^{2+\frac{3v_1}{2}+\frac{v_2}{2}-\frac{v_3}{2}-\frac{3v_4}{2}} y_2^{3+\frac{v_1}{2}+v_2-v_3-\frac{v_4}{2}} y_3^{3-\frac{v_1}{2}-v_2+v_3+\frac{v_4}{2}} y_4^{2-\frac{3v_1}{2}-\frac{v_2}{2}+\frac{v_3}{2}+\frac{3v_4}{2}} \\
& \int_0^\infty \int_0^\infty \int_0^\infty \left( \int_0^\infty K\left[\frac{1}{2}(-2+5v_2+5v_3), \frac{2\pi y_3 \sqrt{1+\frac{1}{u[1]^2}} u[2]}{u[3]} \right] K\left[\frac{1}{2}(-2+5v_2+5v_3), \frac{2\pi y_2 \sqrt{1+u[1]^2} u[3]}{u[4]} \right] u[1]^{-1+\frac{5v_2}{2}-\frac{5v_3}{2}} du[1] \right) \\
& K\left[\frac{1}{2}(-4+5v_1+5v_2+5v_3+5v_4), 2\pi y_4 \sqrt{1+\frac{1}{u[2]^2}} \right] \\
& K\left[\frac{1}{2}(-4+5v_1+5v_2+5v_3+5v_4), 2\pi y_3 \sqrt{(1+u[2]^2) \left(1+\frac{1}{u[3]^2}\right)} \right] \\
& K\left[\frac{1}{2}(-4+5v_1+5v_2+5v_3+5v_4), 2\pi y_2 \sqrt{(1+u[3]^2) \left(1+\frac{1}{u[4]^2}\right)} \right] \\
& K\left[\frac{1}{2}(-4+5v_1+5v_2+5v_3+5v_4), 2\pi y_1 \sqrt{1+u[4]^2} \right] \\
& \left. u[3]^{\frac{1}{2}(-2+5v_1+5v_2-5v_3-5v_4)} (u[2] u[4])^{\frac{1}{2}(-2+5v_1-5v_4)} du[2] du[3] du[4] \right) / \\
& \left( \text{Gamma}\left[\frac{5v_1}{2}\right] \text{Gamma}\left[\frac{5v_2}{2}\right] \text{Gamma}\left[\frac{1}{2}(-1+5v_1+5v_2)\right] \right. \\
& \text{Gamma}\left[\frac{5v_3}{2}\right] \\
& \text{Gamma}\left[\frac{1}{2}(-1+5v_2+5v_3)\right] \\
& \text{Gamma}\left[\frac{1}{2}(-2+5v_1+5v_2+5v_3)\right] \\
& \text{Gamma}\left[\frac{5v_4}{2}\right] \\
& \text{Gamma}\left[\frac{1}{2}(-1+5v_3+5v_4)\right] \\
& \text{Gamma}\left[\frac{1}{2}(-2+5v_2+5v_3+5v_4)\right] \\
& \left. \text{Gamma}\left[\frac{1}{2}(-3+5v_1+5v_2+5v_3+5v_4)\right] \right)
\end{aligned}$$

**FIG. 3.** The  $GL(n)$ pack Whittaker function in dimension 5.

Whittaker function  $W_S^*$ . The timing is from the Mathematica Timingfunction. The results were as follows:

Dimension	v	y	value	timing
3	{5/4, 1/4}	{1, 1}	$2.255480212 \times 10^{-8}$	0.562s
3	{7/6, 5/12}	{1, 1}	$2.255480211 \times 10^{-8}$	
4	{199/520, 23/80, 67/620}	{1, 1, 1}	$1.0910 \times 10^{-15}$	25703.7s
4	{437/1040, 67/260, 213/1040}	{1, 1, 1}	$1.0915 \times 10^{-15}$	
5	{3433/6630, 47/221, 89/510, 3/10}	{1, 1, 1, 1}	$5.1976 \times 10^{-28}$	759.8s
5	{558/1105, 3/10, 22/195, 47/221}	{1, 1, 1, 1}	$5.1972 \times 10^{-28}$	

Given Theorem 2.1, the uniform nature of the periodicity of the integrand should make the application of modern lattice rule techniques [10] practical for direct numerical evaluation of Jacquet's integral. However given the unbounded domain and slow convergence of the integrand this will require considerable adaptation and analysis.

## 9. ACKNOWLEDGEMENTS

The assistance of Eric Stade, Dorian Goldfeld, Ross Barnett and Stephen Joe in the development of this material is gratefully acknowledged.

## REFERENCES

1. P. A. Becker, *On the integration of products of Whittaker functions with respect to the second index*, J. Math. Phys. **45** (2004) p761-773.
2. K. Broughan, *The GL(n)pack manual*. Appendix to *Automorphic forms and L-functions for the group GL(n,R)*. CUP. (to appear).
3. S. Friedberg, *Poincaré Series for GL(n): Fourier expansion, Kloosterman sums and algebro-geometric estimates*. Math. Z. **196** (1987), p165-188.
4. D. Goldfeld, *Automorphic forms and L-functions for the group GL(n,R)*. CUP. (to appear).
5. I. S. Gradshteyn and I. M. Ryzhik, *Tables of integrals series and products.*, Academic Press, 6th ed., 2000.
6. T. Ishii, *A remark on Whittaker functions on SL(n,R)*, Ann.Inst.Fourier.Grenoble **55** (2005), p483-492.
7. H. Jacquet, *Fonctions de Whittaker associées aux groupes de Chevalley*, Bull. Soc. Math. France **95** (1967), p243-309.
8. N. N. Lebedev, *Special functions and their applications*, Dover, 1972.
9. F. W. J. Olver, *Asymptotics and Special Functions*, A. K. Peters, 1997.
10. I. H. Sloan and T. R. Osborn, *Multiple integration over bounded and unbounded regions.*, J. Comp. App. Math. **17** (1987), p181-196.
11. E. Stade, *On explicit integral formulas for GL(n,R)-Whittaker Functions.*, Duke Math. J. **60** (1990), p313-362.
12. E. Stade, *Mellin transforms of GL(n,R) Whittaker functions.* Amer. J. Math. **123** (2001), p121-161 .
13. E. Stade, *Archimedean L-factors on GL(n)x GL(n) and generalized Barnes integrals*, Israel J. Math. **127** (2002), p201-219.
14. A. H. Stroud, *Approximate Calculation of Multiple Integrals*. Prentice-Hall, 1971.
15. A. Terras, *Harmonic analysis on symmetric spaces and applications I*. Springer-Verlag, 1985.
16. E. T. Whittaker and G. N. Watson, *A course in modern analysis*. 4th ed. CUP, 1963.